



# Free Home Delivery

(Online mail order pharmacy)

Enrolling in home delivery is easy with  
this simple, 2-step process

**1** **Create an account online** at <https://ridgeway.pharmacy/Create%20Account> or simply call Ridgeway Mail Order Pharmacy at 1-800-630-3214 and press 0 to speak with a member of the pharmacy staff who will help you. The hours of operation are Monday–Friday, 7:00 AM–3:00 PM ET.

**2** **Contact your doctor who prescribed CONTRAVE** to let them know you've enrolled in the Home Delivery program and ask them to complete the process in 1 of 3 ways:

**EMR:** They select “Ridgeway Mail Order Pharmacy” (2824 US Hwy 93 North, Victor, MT 59875) as the pharmacy of choice (NCPDP#: 2706488) and submit the prescription

**Phone:** They call Ridgeway Mail Order Pharmacy at 1-800-630-3214 and press 2 to authorize the prescription

**Fax:** They fax your prescription to Ridgeway Mail Order Pharmacy at 1-406-642-6050

**Note:** If you currently take CONTRAVE and it is covered by your insurance, you'll pay \$99 per month by switching to the free Home Delivery program.

CONTRAVE® (naltrexone HCl/bupropion HCl) is a prescription weight-loss medicine that may help adults with obesity (BMI greater than or equal to 30 kg/m<sup>2</sup>), or are overweight (BMI greater than or equal to 27 kg/m<sup>2</sup>) with at least one weight-related medical condition, lose weight and keep the weight off. CONTRAVE should be used along with diet and exercise.

Please see Important Safety Information for CONTRAVE on the next page.

## Important Safety Information

One of the ingredients in CONTRAVE, bupropion, may increase the risk of suicidal thinking in children, adolescents, and young adults. CONTRAVE patients should be monitored for suicidal thoughts and behaviors. In patients taking bupropion for smoking cessation, serious neuropsychiatric adverse events have been reported. CONTRAVE is not approved for children under 18.

Stop taking CONTRAVE and call your healthcare provider right away if you experience thoughts about suicide or dying; depression, or anxiety; panic attacks; trouble sleeping; irritability; aggression; mania; or other unusual changes in behavior or mood.

Do not take CONTRAVE if you: have uncontrolled hypertension; have or have had seizures or an eating disorder; use other medicines that contain bupropion; are dependent on opioid pain medicines, use medicines to help stop taking opioids, or are in opiate withdrawal; drink a lot of alcohol and abruptly stop drinking, or take sedatives, benzodiazepines, or anti-seizure medicines and you abruptly stop using them; or are taking monoamine oxidase inhibitors (MAOIs). Do not start CONTRAVE until you have stopped taking your MAOI for at least 14 days. Do not take CONTRAVE if you are allergic to any of the ingredients in CONTRAVE. Do not take CONTRAVE if you are pregnant or planning to become pregnant or are breastfeeding.

Before taking CONTRAVE, tell your healthcare provider about all of your current or past health conditions.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Do not take any other medicines while you are taking CONTRAVE unless your healthcare provider says it is okay.

If you have a seizure while taking CONTRAVE, stop taking CONTRAVE and call your healthcare provider right away.

Additional serious side effects may include: opioid overdose or sudden opioid withdrawal; severe allergic reactions; increases in blood pressure or heart rate; liver damage or hepatitis; manic episodes; visual problems (glaucoma); and increased risk of low blood sugar (hypoglycemia) in people with type 2 diabetes mellitus who take certain medicines to treat their diabetes.

The most common side effects of CONTRAVE include nausea, constipation, headache, vomiting, dizziness, trouble sleeping, dry mouth, and diarrhea.

These are not all the possible side effects of CONTRAVE. Please refer to the full Prescribing Information, including Medication Guide, for CONTRAVE at [www.CONTRAVE.com](http://www.CONTRAVE.com) or talk to your doctor.

You are encouraged to report negative side effects of drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.



OREXIGEN®

CONTRAVE® is a registered trademark of Orexigen Therapeutics, Inc.  
©2017 Orexigen Therapeutics, Inc. CON-0417 05/17



**Contrace**®  
(naltrexone HCl/bupropion HCl)  
8 mg/90 mg • Extended-Release Tablets